

Misericordia University

## Misericordia Digital Commons

---

Student Research Poster Presentations 2021

Student Research Poster Presentations

---

2021

### Reducing Dose In Coronary Computed Tomography Angiography (CCTA)

Mariah Ambrosio  
ambrosim@misericordia.edu

Follow this and additional works at: [https://digitalcommons.misericordia.edu/research\\_posters2021](https://digitalcommons.misericordia.edu/research_posters2021)



Part of the [Medicine and Health Sciences Commons](#)

---

#### Recommended Citation

Ambrosio, Mariah, "Reducing Dose In Coronary Computed Tomography Angiography (CCTA)" (2021).  
*Student Research Poster Presentations 2021*. 19.  
[https://digitalcommons.misericordia.edu/research\\_posters2021/19](https://digitalcommons.misericordia.edu/research_posters2021/19)

This Poster is brought to you for free and open access by the Student Research Poster Presentations at Misericordia Digital Commons. It has been accepted for inclusion in Student Research Poster Presentations 2021 by an authorized administrator of Misericordia Digital Commons. For more information, please contact [jluksa@misericordia.edu](mailto:jluksa@misericordia.edu), [mcech@misericordia.edu](mailto:mcech@misericordia.edu).



## What is CT?

- Computed tomography (CT) is the mechanism of creating cross-sectional tomographic planes of any part of the body
- Cross-sectional images generated can be reformatted in coronal, sagittal and axial planes

(Long, Rollins & Smith, 2016, p. 302)

## What is CCTA?

- Coronary computed tomography angiography (CCTA) is a noninvasive imaging procedure to aid in diagnosis coronary artery disease (CAD)
- Appropriate for diagnosis and risk assessment in those with low to intermediate CAD risk

(Hamilton-Craig et al., 2020, p. 26)




Image 1: Arrow pointing to calcified plaque in proximal left anterior descending coronary artery

(Renker et al., 2011, p. 394)

## Dose Received From CCTA Exams

- CCTA studies average 15.7 millisieverts (mSv) of effective dose, some studies reaching up to 20 mSv or more (Hamilton-Craig et al., 2020, p. 26).
- Risk of exposure-induced cancer death (REID) was 1 in 746 men and 1 in 508 women undergoing CCTA exams
- REID values found to be considerably higher in younger women

(Mahmoodi & Chaparian, 2019, p. 1134)

## Electrocardiogram (ECG)-Triggered Prospective Gating

- Associated with lowest-dose CCTA scans
- Monitors patient's heart rate and scans during most useful parts of the cardiac cycle
  - Typically time elapsed between two successive R waves of QRS signal on ECG; R-R interval
- Prospective gating activates tube to scan only during predetermined points in the cardiac cycle
- Accounts for an approximate 67% to 72% decrease in radiation dose during CCTA exams

(Corbett, 2020, p. 405)




Image 2: Radiation exposure during retrospective gating (tube active throughout cardiac cycle) and prospective gating

(Hamilton-Craig et al., 2020, p. 27)

## Proper Use & Limitations of Prospective Gating

- Effective prospective gating requires regular, slow heart rate, typically 50-65 beats per minute (BPM)
- Patients whose heart rate exceeds 70 bpm do not qualify for prospective gating
  - Increased risk of motion artifact, resulting in repeating entire scan & increased patient dose

(Fornell, 2019, p. 27)




Image 3: Effect of heart rate on prospectively gated CCTA scan

(Hamilton-Craig et al., 2019, p. 28)

## Reducing Tube Potential

- Studies have shown the most important factor of controlling dose is adjustment of x-ray tube voltage
- Due to contrast administration, typical tube voltage for CCTA exams around 120 kilovolts peak (kVp)
- Reducing tube potential from the standard 120 kVp to 80-100 kVp significantly reduced dose approximately 31% to 40%

(Fornell, 2019, p. 26; Hamilton-Craig et al., 2020, p. 31)




Image 4: Recommended tube voltage for patient's BMI

(Hamilton-Craig et al., 2020, p. 31)

## Proper Use & Limitations of Tube Potential Reduction

- Lowering kVp lowers radiation dose while simultaneously increases image noise
- Selecting tube potential involves a trade-off between image noise and dose
- Better suited for use on average-sized or smaller patients
  - Body mass index (BMI) < 30 kg/m<sup>2</sup>
- Reducing tube potential on obese patients results in scans with inadequate image quality

(Corbett, 2020, p. 405; Fornell, 2019, p. 25)

## Bolus Tracking

- Begins with a delay to allow contrast to get to target area, takes intermittent pictures of descending aorta until it reaches threshold of 120 Hounsfield Units (HU)
- Rapid contrast injection (4-6ml/sec) required in CCTA to highlight the arteries and vessels
- Bolus tracking (BT) used to estimate contrast arrival timing adding to approximately 12% of total dose
- Delay of 15 seconds and intermittent scans of 1-2s showed a significant reduction in dose without interfering with image quality

(Nishimura Matsumoto et al., 2018, pp. 313-314)




Image 5: BT effective radiation dose among protocols

(Nishimura Matsumoto et al., 2018, p. 314)

## Conclusion

- CCTA is becoming an increasingly valuable diagnostic tool in diagnosing CAD
- Dose reduction techniques implemented during CCTA exams provide diagnostic quality scans while lowering dose to the patient
- Each method can be used independently or in conjunction with another
- Understanding the limitations and drawbacks of each method is extremely important




Image 6: Relationship between dose and proportion of patients receiving one or more reduction method

(Hamilton-Craig et al., 2020, p. 28)